

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
30 January 2003 (30.01.2003)

PCT

(10) International Publication Number  
WO 03/008613 A2(51) International Patent Classification<sup>7</sup>: C12P 13/00

(21) International Application Number: PCT/EP02/07373

(22) International Filing Date: 3 July 2002 (03.07.2002)


(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
101 35 053.8 18 July 2001 (18.07.2001) DE  
60/306,869 23 July 2001 (23.07.2001) USCZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VN, YU, ZA, ZM, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,  
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).(71) Applicant (for all designated States except US): DE-  
GUSSA AG [DE/DE]; Bennigsenplatz 1, 40474 Düssel-  
dorf (DE).Declaration under Rule 4.17:  
— of inventorship (Rule 4.17(iv)) for US only

(72) Inventor; and

Published:

(75) Inventor/Applicant (for US only): RIEPING, Mechthild  
[DE/DE]; Mönkebergstrasse 1, 33619 Bielefeld (DE).— without international search report and to be republished  
upon receipt of that report(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.(54) Title: PROCESS FOR THE PREPARATION OF L-AMINO ACIDS USING STRAINS OF THE ENTEROBACTERIACEAE  
FAMILY WHICH CONTAIN AN ENHANCED SODA GENE(57) Abstract: The invention relates to a process for the preparation of L-amino acids, in particular L-threonine, in which the fol-  
lowing steps are carried out: a) fermentation of microorganisms of the Enterobacteriaceae family which produce the desired L-amino  
acid and in which at least the sodA gene, or the nucleotide sequence which codes for this, is enhanced, in particular over-expressed,  
b) concentration of the desired L-amino acid in the medium or in the cells of the bacteria, and c) isolation of the desired L-amino  
acid.

**Process for the Preparation of L-Amino Acids using  
Strains of the Enterobacteriaceae Family which Contain  
an Enhanced *sodA* Gene**

Field of the Invention

- 5 This invention relates to a process for the fermentative preparation of L-amino acids, in particular L-threonine, using strains of the Enterobacteriaceae family in which at least the *sodA* gene is enhanced.

Prior Art

- 10 L-Amino acids, in particular L-threonine, are used in human medicine and in the pharmaceuticals industry, in the foodstuffs industry and very particularly in animal nutrition.

It is known to prepare L-amino acids by fermentation of  
15 strains of Enterobacteriaceae, in particular *Escherichia coli* (*E. coli*) and *Serratia marcescens*. Because of their great importance, work is constantly being undertaken to improve the preparation processes. Improvements to the process can relate to fermentation measures, such as e.g.  
20 stirring and supply of oxygen, or the composition of the nutrient media, such as e.g. the sugar concentration during the fermentation, or the working up to the product form, by e.g. ion exchange chromatography, or the intrinsic output properties of the microorganism itself.

- 25 Methods of mutagenesis, selection and mutant selection are used to improve the output properties of these microorganisms. Strains which are resistant to antimetabolites, such as e.g. the threonine analogue  $\alpha$ -amino- $\beta$ -hydroxyvaleric acid (AHV), or are auxotrophic for  
30 metabolites of regulatory importance and produce L-amino acid, such as e.g. L-threonine, are obtained in this manner.

Methods of the recombinant DNA technique have also been employed for some years for improving the strain of strains of the Enterobacteriaceae family which produce L-amino acids, by amplifying individual amino acid biosynthesis genes and investigating the effect on the production.

#### Object of the Invention

The object of the invention is to provide new measures for improved fermentative preparation of L-amino acids, in particular L-threonine.

#### Summary of the Invention

The invention provides a process for the fermentative preparation of L-amino acids, in particular L-threonine, using microorganisms of the Enterobacteriaceae family which in particular already produce L-amino acids and in which at least the nucleotide sequence which codes for the *sodA* gene is enhanced.

#### Detailed Description of the Invention

Where L-amino acids or amino acids are mentioned in the following, this means one or more amino acids, including their salts, chosen from the group consisting of L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine. L-Threonine is particularly preferred.

The term "enhancement" in this connection describes the increase in the intracellular activity of one or more enzymes or proteins in a microorganism which are coded by the corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or a gene or allele which codes for a corresponding enzyme or

protein with a high activity, and optionally combining these measures.

By enhancement measures, in particular over-expression, the activity or concentration of the corresponding protein is  
5 in general increased by at least 10%, 25%, 50%, 75%, 100%, 150%, 200%, 300%, 400% or 500%, up to a maximum of 1000% or 2000%, based on that of the wild-type protein or the activity or concentration of the protein in the starting microorganism.

10 The process comprises carrying out the following steps:

- a) fermentation of microorganisms of the Enterobacteriaceae family in which at least the *sodA* gene is enhanced,
- 15 b) concentration of the corresponding L-amino acid in the medium or in the cells of the microorganisms of the Enterobacteriaceae family, and
- c) isolation of the desired L-amino acid, constituents of the fermentation broth and/or the biomass in its entirety or portions (> 0 to 100 %) thereof  
20 optionally remaining in the product.

The microorganisms which the present invention provides can produce L-amino acids from glucose, sucrose, lactose, fructose, maltose, molasses, optionally starch, optionally cellulose or from glycerol and ethanol. They are  
25 representatives of the Enterobacteriaceae family chosen from the genera *Escherichia*, *Erwinia*, *Providencia* and *Serratia*. The genera *Escherichia* and *Serratia* are preferred. Of the genus *Escherichia* the species *Escherichia coli* and of the genus *Serratia* the species *Serratia*  
30 *marcescens* are to be mentioned in particular.

Suitable strains, which produce L-threonine in particular, of the genus *Escherichia*, in particular of the species *Escherichia coli*, are, for example

- Escherichia coli TF427
- 5 Escherichia coli H4578
- Escherichia coli KY10935
- Escherichia coli VNIIGenetika MG442
- Escherichia coli VNIIGenetika M1
- Escherichia coli VNIIGenetika 472T23
- 10 Escherichia coli BKIIM B-3996
- Escherichia coli kat 13
- Escherichia coli KCCM-10132.

Suitable L-threonine-producing strains of the genus *Serratia*, in particular of the species *Serratia marcescens*,  
15 are, for example

- Serratia marcescens* HNr21
- Serratia marcescens* TLr156
- Serratia marcescens* T2000.

Strains from the Enterobacteriaceae family which produce L-  
20 threonine preferably have, inter alia, one or more genetic or phenotypic features chosen from the group consisting of: resistance to  $\alpha$ -amino- $\beta$ -hydroxyvaleric acid, resistance to thialysine, resistance to ethionine, resistance to  $\alpha$ -methylserine, resistance to diaminosuccinic acid,  
25 resistance to  $\alpha$ -aminobutyric acid, resistance to borrelidin, resistance to rifampicin, resistance to valine analogues, such as, for example, valine hydroxamate, resistance to purine analogues, such as, for example, 6-dimethylaminopurine, a need for L-methionine, optionally a  
30 partial and compensatable need for L-isoleucine, a need for meso-diaminopimelic acid, auxotrophy in respect of threonine-containing dipeptides, resistance to L-threonine, resistance to L-homoserine, resistance to L-lysine, resistance to L-methionine, resistance to L-glutamic acid,

resistance to L-aspartate, resistance to L-leucine,  
resistance to L-phenylalanine, resistance to L-serine,  
resistance to L-cysteine, resistance to L-valine,  
sensitivity to fluoropyruvate, defective threonine  
5 dehydrogenase, optionally an ability for sucrose  
utilization, enhancement of the threonine operon,  
enhancement of homoserine dehydrogenase I-aspartate kinase  
I, preferably of the feed back resistant form, enhancement  
of homoserine kinase, enhancement of threonine synthase,  
10 enhancement of aspartate kinase, optionally of the feed  
back resistant form, enhancement of aspartate semialdehyde  
dehydrogenase, enhancement of phosphoenol pyruvate  
carboxylase, optionally of the feed back resistant form,  
enhancement of phosphoenol pyruvate synthase, enhancement  
15 of transhydrogenase, enhancement of the RhtB gene product,  
enhancement of the RhtC gene product, enhancement of the  
YfiK gene product, enhancement of a pyruvate carboxylase,  
and attenuation of acetic acid formation.

It has been found that microorganisms of the  
20 Enterobacteriaceae family produce L-amino acids, in  
particular L-threonine, in an improved manner after  
enhancement, in particular over-expression, of the *sodA*  
gene.

The nucleotide sequences of the genes of *Escherichia coli*  
25 belong to the prior art and can also be found in the genome  
sequence of *Escherichia coli* published by Blattner et al.  
(*Science* 277: 1453-1462 (1997)).

The following information, inter alia, on the *sodA* gene is  
known from the prior art:

30 Description: Superoxide dismutase  
EC No. 1.15.1.1

- Reference: Touati; Journal of Bacteriology 155(3):  
1078-1087 (1983); Schrum and Hassan; Free  
Radical Biology and Medicine 17(3): 209-213  
(1994); Benov and Fridovich; Archives of  
5 Biochemistry and Biophysics 322(1): 291-294  
(1995)

Accession No.: AE000465

The nucleic acid sequences can be found in the databanks of  
the National Center for Biotechnology Information (NCBI) of  
10 the National Library of Medicine (Bethesda, MD, USA), the  
nucleotide sequence databank of the European Molecular  
Biologies Laboratories (EMBL, Heidelberg, Germany or  
Cambridge, UK) or the DNA databank of Japan (DDBJ, Mishima,  
Japan).

15 The genes described in the text references mentioned can be  
used according to the invention. Alleles of the genes which  
result from the degeneracy of the genetic code or due to  
"sense mutations" of neutral function can furthermore be  
used.

20 To achieve an enhancement, for example, expression of the  
genes or the catalytic properties of the proteins can be  
increased. The two measures can optionally be combined.

To achieve an over-expression, the number of copies of the  
corresponding genes can be increased, or the promoter and  
25 regulation region or the ribosome binding site upstream of  
the structural gene can be mutated. Expression cassettes  
which are incorporated upstream of the structural gene act  
in the same way. By inducible promoters, it is  
additionally possible to increase the expression in the  
30 course of fermentative L-threonine production. The  
expression is likewise improved by measures to prolong the  
life of the m-RNA. Furthermore, the enzyme activity is  
also increased by preventing the degradation of the enzyme  
protein. The genes or gene constructs can either be

present in plasmids with a varying number of copies, or can be integrated and amplified in the chromosome.

Alternatively, an over-expression of the genes in question can furthermore be achieved by changing the composition of  
5 the media and the culture procedure.

Instructions in this context can be found by the expert, inter alia, in Chang and Cohen (Journal of Bacteriology 134: 1141-1156 (1978)), in Hartley and Gregori (Gene 13: 347-353 (1981)), in Amann and Brosius (Gene 40: 183-190  
10 (1985)), in de Broer et al. (Proceedings of the National Academy of Sciences of the United States of America 80: 21-25 (1983)), in LaVallie et al. (BIO/TECHNOLOGY 11: 187-193 (1993)), in PCT/US97/13359, in Llosa et al. (Plasmid 26: 222-224 (1991)), in Quandt and Klipp (Gene 80: 161-169  
15 (1989)), in Hamilton et al. (Journal of Bacteriology 171: 4617-4622 (1989)), in Jensen and Hammer (Biotechnology and Bioengineering 58: 191-195 (1998)) and in known textbooks of genetics and molecular biology.

Plasmid vectors which are capable of replication in  
20 Enterobacteriaceae, such as e.g. cloning vectors derived from pACYC184 (Bartolomé et al.; Gene 102: 75-78 (1991)), pTrc99A (Amann et al.; (Gene 69: 301-315 (1988)) or pSC101 derivatives (Vočke and Bastia; Proceedings of the National Academy of Sciences of the United States of America 80  
25 (21): 6557-6561 (1983)) can be used. A strain transformed with a plasmid vector, where the plasmid vector carries at least the sodA gene, or the nucleotide sequence which codes for this, can be employed in a process according to the invention.

30 It is also possible to transfer mutations which affect the expression of the particular gene into various strains by sequence exchange (Hamilton et al.; Journal of Bacteriology 171: 4617-4622 (1989)), conjugation or transduction.



It may furthermore be advantageous for the production of L-amino acids, in particular L-threonine, with strains of the Enterobacteriaceae family, in addition to enhancement of the *sodA* gene, for one or more enzymes of the known  
5 threonine biosynthesis pathway or enzymes of anaplerotic metabolism or enzymes for the production of reduced nicotinamide adenine dinucleotide phosphate or enzymes of glycolysis or PTS enzymes or enzymes of sulfur metabolism to be enhanced.

10 Thus, for example, at the same time one or more of the genes chosen from the group consisting of

- the *thrABC* operon which codes for aspartate kinase, homoserine dehydrogenase, homoserine kinase and threonine synthase (US-A-4,278,765),
- 15 • the *pyc* gene of *Corynebacterium glutamicum* which codes for pyruvate carboxylase (WO 99/18228),
- the *pps* gene which codes for phosphoenol pyruvate synthase (Molecular and General Genetics 231(2): 332-336 (1992)),
- 20 • the *ppc* gene which codes for phosphoenol pyruvate carboxylase (Gene 31: 279-283 (1984)),
- the *pntA* and *pntB* genes which code for transhydrogenase (European Journal of Biochemistry 158: 647-653 (1986)),
- the *rhtB* gene which imparts homoserine resistance (EP-A-  
25 0 994 190),
- the *mgo* gene which codes for malate:quinone oxidoreductase (WO 02/06459),
- the *rhtC* gene which imparts threonine resistance (EP-A-1  
013 765),

- the thrE gene of Corynebacterium glutamicum which codes for the threonine export protein (WO 01/92545),
- the gdhA gene which codes for glutamate dehydrogenase (Nucleic Acids Research 11: 5257-5266 (1983); Gene 23: 199-209 (1983)),
- the hns gene which codes for the DNA-binding protein HLP-II (Molecular and General Genetics 212: 199-202 (1988)),
- the pgm gene which codes for phosphoglucomutase (Journal of Bacteriology 176: 5847-5851 (1994)),
- the fba gene which codes for fructose biphosphate aldolase (Biochemical Journal 257: 529-534 (1989)),
- the ptsH gene of the ptsHIcrr operon which codes for the phosphohistidine protein hexose phosphotransferase of the phosphotransferase system PTS (Journal of Biological Chemistry 262: 16241-16253 (1987)),
- the ptsI gene of the ptsHIcrr operon which codes for enzyme I of the phosphotransferase system PTS (Journal of Biological Chemistry 262: 16241-16253 (1987)),
- the crr gene of the ptsHIcrr operon which codes for the glucose-specific IIA component of the phosphotransferase system PTS (Journal of Biological Chemistry 262: 16241-16253 (1987)),
- the ptsG gene which codes for the glucose-specific IIBC component (Journal of Biological Chemistry 261: 16398-16403 (1986)),
- the lrp gene which codes for the regulator of the leucine regulon (Journal of Biological Chemistry 266: 10768-10774 (1991)).

- the mopB gene which codes for 10 Kd chaperone (Journal of Biological Chemistry 261: 12414-12419 (1986)) and is also known by the name groES,
- 5 • the ahpC gene of the ahpCF operon which codes for the small sub-unit of alkyl hydroperoxide reductase (Proceedings of the National Academy of Sciences of the United States of America 92: 7617-7621 (1995)),
- 10 • the ahpF gene of the ahpCF operon which codes for the large sub-unit of alkyl hydroperoxide reductase (Proceedings of the National Academy of Sciences of the United States of America 92: 7617-7621 (1995)),
- the cysK gene which codes for cysteine synthase A (Journal of Bacteriology 170: 3150-3157 (1988)),
- 15 • the cysB gene which codes for the regulator of the cys regulon (Journal of Biological Chemistry 262: 5999-6005 (1987)),
- 20 • the cysJ gene of the cysJIH operon which codes for the flavoprotein of NADPH sulfite reductase (Journal of Biological Chemistry 264: 15796-15808 (1989), Journal of Biological Chemistry 264: 15726-15737 (1989)),
- the cysI gene of the cysJIH operon which codes for the haemoprotein of NADPH sulfite reductase (Journal of Biological Chemistry 264: 15796-15808 (1989), Journal of Biological Chemistry 264: 15726-15737 (1989)),
- 25 • the cysH gene of the cysJIH operon which codes for adenylyl sulfate reductase (Journal of Biological Chemistry 264: 15796-15808 (1989), Journal of Biological Chemistry 264: 15726-15737 (1989)),
- 30 • the phoB gene of the phoBR operon which codes for the positive regulator PhoB of the pho regulon (Journal of Molecular Biology 190 (1): 37-44 (1986)),

- the phoR gene of the phoBR operon which codes for the sensor protein of the pho regulon (Journal of Molecular Biology 192 (3): 549-556 (1986)),
- the phoE gene which codes for protein E of the outer  
5 cell membrane (Journal of Molecular Biology 163 (4): 513-532 (1983)),
- the malE gene which codes for the periplasmic binding protein of maltose transport (Journal of Biological Chemistry 259 (16): 10606-10613 (1984)),
- 10 • the pykF gene which codes for fructose-stimulated pyruvate kinase I (Journal of Bacteriology 177 (19): 5719-5722 (1995)),
- the talB gene which codes for transaldolase B (Journal of Bacteriology 177 (20): 5930-5936 (1995)),
- 15 • the rseA gene of the rseABC operon which codes for a membrane protein with anti-sigmaE activity (Molecular Microbiology 24 (2): 355-371 (1997)),
- the rseC gene of the rseABC operon which codes for a global regulator of the sigmaE factor (Molecular  
20 Microbiology 24 (2): 355-371 (1997)),
- the pfkB gene which codes for 6-phosphofructokinase II (Gene 28 (3): 337-342 (1984)),
- the sucA gene of the sucABCD operon which codes for the decarboxylase sub-unit of 2-ketoglutarate dehydrogenase  
25 (European Journal of Biochemistry 141 (2): 351-359 (1984)),
- the sucB gene of the sucABCD operon which codes for the dihydrolipoyltranssuccinase E2 sub-unit of 2-ketoglutarate dehydrogenase (European Journal of  
30 Biochemistry 141 (2): 351-359 (1984)),

- the *sucC* gene of the *sucABCD* operon which codes for the  $\beta$ -sub-unit of succinyl-CoA synthetase (Biochemistry 24 (22): 6245-6252 (1985)) and
- the *sucD* gene of the *sucABCD* operon which codes for the  $\alpha$ -sub-unit of succinyl-CoA synthetase (Biochemistry 24 (22): 6245-6252 (1985)),

can be enhanced, in particular over-expressed.

It may furthermore be advantageous for the production of L-amino acids, in particular L-threonine, in addition to enhancement of the *sodA* gene, for one or more of the genes chosen from the group consisting of

- the *tdh* gene which codes for threonine dehydrogenase (Journal of Bacteriology 169: 4716-4721 (1987)),
- the *mdh* gene which codes for malate dehydrogenase (E.C. 1.1.1.37) (Archives in Microbiology 149: 36-42 (1987)),
- the gene product of the open reading frame (orf) *yjfa* (Accession Number AAC77180 of the National Center for Biotechnology Information (NCBI, Bethesda, MD, USA)),
- the gene product of the open reading frame (orf) *ytfp* (Accession Number AAC77179 of the National Center for Biotechnology Information (NCBI, Bethesda, MD, USA)),
- the *pckA* gene which codes for the enzyme phosphoenolpyruvate carboxykinase (Journal of Bacteriology 172: 7151-7156 (1990)),
- the *poxB* gene which codes for pyruvate oxidase (Nucleic Acids Research 14(13): 5449-5460 (1986)),
- the *aceA* gene which codes for the enzyme isocitrate lyase (Journal of Bacteriology 170: 4528-4536 (1988)),

- the dgsA gene which codes for the DgsA regulator of the phosphotransferase system (Bioscience, Biotechnology and Biochemistry 59: 256-251 (1995)) and is also known under the name of the mlc gene,
- 5 • the fruR gene which codes for the fructose repressor (Molecular and General Genetics 226: 332-336 (1991)) and is also known under the name of the cra gene and
- the rpoS gene which codes for the sigma<sup>38</sup> factor (WO 01/05939) and is also known under the name of the katF  
10 gene,

to be attenuated, in particular eliminated or for the expression thereof to be reduced.

The term "attenuation" in this connection describes the reduction or elimination of the intracellular activity of  
15 one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by using a weak promoter or a gene or allele which codes for a corresponding enzyme with a low activity or inactivates the corresponding enzyme (protein) or gene, and optionally  
20 combining these measures.

By attenuation measures, the activity or concentration of the corresponding protein is in general reduced to 0 to 75%, 0 to 50%, 0 to 25%, 0 to 10% or 0 to 5% of the activity or concentration of the wild-type protein or of  
25 the activity or concentration of the protein in the starting microorganism.

It may furthermore be advantageous for the production of L-amino acids, in particular L-threonine, in addition to enhancement of the sodA gene, to eliminate undesirable side  
30 reactions (Nakayama: "Breeding of Amino Acid Producing Microorganisms", in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

The microorganisms produced according to the invention can be cultured in the batch process (batch culture), the fed batch process (feed process) or the repeated fed batch process (repetitive feed process). A summary of known  
5 culture methods is described in the textbook by Chmiel (Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik [Bioprocess Technology 1. Introduction to Bioprocess Technology (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und  
10 periphere Einrichtungen [Bioreactors and Peripheral Equipment] (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained  
15 in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981).

Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and optionally  
20 cellulose, oils and fats, such as e.g. soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as e.g. palmitic acid, stearic acid and linoleic acid, alcohols, such as e.g. glycerol and ethanol, and organic acids, such as e.g. acetic acid, can be used as the source  
25 of carbon. These substances can be used individually or as a mixture.

Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds,  
30 such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate, can be used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture.

Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of  
5 metals, such as e.g. magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the abovementioned substances. Suitable precursors can moreover be added to the culture  
10 medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds,  
15 such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH of the culture. Antifoams, such as e.g. fatty acid polyglycol esters, can be employed to control the development of foam. Suitable substances having a selective action, e.g. antibiotics, can  
20 be added to the medium to maintain the stability of plasmids. To maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such as e.g. air, are introduced into the culture. The temperature of the culture is usually 25°C to 45°C, and preferably 30°C to  
25 40°C. Culturing is continued until a maximum of L-amino acids or L-threonine has formed. This target is usually reached within 10 hours to 160 hours.

The analysis of L-amino acids can be carried out by anion exchange chromatography with subsequent ninhydrin  
30 derivation, as described by Spackman et al. (Analytical Chemistry 30: 1190-1206 (1958)), or it can take place by reversed phase HPLC as described by Lindroth et al. (Analytical Chemistry 51: 1167-1174 (1979)).

The process according to the invention is used for the  
35 fermentative preparation of L-amino acids, such as, for



example, L-threonine, L-isoleucine, L-valine, L-methionine, L-homoserine and L-lysine, in particular L-threonine.

The present invention is explained in more detail in the following with the aid of embodiment examples.

- 5 The minimal (M9) and complete media (LB) for Escherichia coli used are described by J.H. Miller (A Short Course in Bacterial Genetics (1992), Cold Spring Harbor Laboratory Press). The isolation of plasmid DNA from Escherichia coli and all techniques of restriction, ligation, Klenow and  
10 alkaline phosphatase treatment are carried out by the method of Sambrook et al. (Molecular Cloning - A Laboratory Manual (1989) Cold Spring Harbor Laboratory Press). Unless described otherwise, the transformation of Escherichia coli is carried out by the method of Chung et al. (Proceedings  
15 of the National Academy of Sciences of the United States of America 86: 2172-2175 (1989)).

The incubation temperature for the preparation of strains and transformants is 37°C.

#### Example 1

- 20 Construction of the expression plasmid pTrc99AsodA

The sodA gene from E. coli K12 is amplified using the polymerase chain reaction (PCR) and synthetic oligonucleotides. Starting from the nucleotide sequence of the sodA gene in E. coli K12 MG1655 (Accession Number  
25 AE000465, Blattner et al. (Science 277: 1453-1462 (1997))), PCR primers are synthesized (MWG Biotech, Ebersberg, Germany). The sequences of the primers are modified such that recognition sites for restriction enzymes are formed. The recognition sequence for XbaI is chosen for the sodA1  
30 primer and the recognition sequence for HindIII for the sodA2 primer, which are marked by underlining in the nucleotide sequence shown below:

sodA1: 5' - CGCGGCATCTAGAAATCTGACGCCCCGACAATAC - 3'  
(SEQ ID No. 1)

sodA2: 5' - CGCCTCATTGAAGCTTGCGGCAAATG - 3'  
(SEQ ID No. 2)

5 The chromosomal E. coli K12 MG1655 DNA employed for the PCR is isolated according to the manufacturer's instructions with "Qiagen Genomic-tips 100/G" (QIAGEN, Hilden, Germany). A DNA fragment approx. 700 bp in size can be amplified with the specific primers under standard PCR conditions (Innis  
10 et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press) with Pfu-DNA polymerase (Promega Corporation, Madison, USA).

The PCR product is ligated according to the manufacturer's instructions with the vector pCR-Blunt II-TOPO (Zero Blunt  
15 TOPO PCR Cloning Kit, Invitrogen, Groningen, The Netherlands) and transformed into the E. coli strain TOP10. Selection of plasmid-carrying cells takes place on LB agar, to which 50 µg/ml kanamycin are added. After isolation of the plasmid DNA, the vector pCR-Blunt II-TOPO-sodA is  
20 cleaved with the restriction enzymes HindIII and XbaI and, after separation in 0.8% agarose gel, the sodA fragment is isolated with the aid of the QIAquick Gel Extraction Kit (QIAGEN, Hilden, Germany). The vector pTrc99A (Pharmacia Biotech, Uppsala, Sweden) is cleaved with the enzymes  
25 HindIII and XbaI and ligation is carried out with the sodA fragment isolated.

The E. coli strain XL1-Blue MRF' (Stratagene, La Jolla, USA) is transformed with the ligation batch and plasmid-carrying cells are selected on LB agar, to which 50 µg/ml  
30 ampicillin are added. Successful cloning can be demonstrated after plasmid DNA isolation by control cleavage with the enzymes BglI, Eco91I and PvuII. The plasmid is sequenced (Figure 2).

Example 2

Preparation of L-threonine with the strain  
MG442/pTrc99AsodA

The L-threonine-producing E. coli strain MG442 is described  
5 in the patent specification US-A- 4,278,765 and deposited  
as CMIM B-1628 at the Russian National Collection for  
Industrial Microorganisms (VKPM, Moscow, Russia).

The strain MG442 is transformed with the expression plasmid  
pTrc99AsodA described in example 1 and with the vector  
10 pTrc99A and plasmid-carrying cells are selected on LB agar  
with 50 µg/ml ampicillin. The strains MG442/pTrc99AsodA and  
MG442/pTrc99A are formed in this manner. Selected  
individual colonies are then multiplied further on minimal  
medium with the following composition: 3.5 g/l Na<sub>2</sub>HPO<sub>4</sub>\*2H<sub>2</sub>O,  
15 1.5 g/l KH<sub>2</sub>PO<sub>4</sub>, 1 g/l NH<sub>4</sub>Cl, 0.1 g/l MgSO<sub>4</sub>\*7H<sub>2</sub>O, 2 g/l  
glucose, 20 g/l agar, 50 mg/l ampicillin. The formation of  
L-threonine is checked in batch cultures of 10 ml contained  
in 100 ml conical flasks. For this, 10 ml of preculture  
medium of the following composition: 2 g/l yeast extract,  
20 10 g/l (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 1 g/l KH<sub>2</sub>PO<sub>4</sub>, 0.5 g/l MgSO<sub>4</sub>\*7H<sub>2</sub>O, 15 g/l  
CaCO<sub>3</sub>, 20 g/l glucose, 50 mg/l ampicillin are inoculated  
and the batch is incubated for 16 hours at 37°C and 180 rpm  
on an ESR incubator from Kühner AG (Birsfelden,  
Switzerland).

25 250 µl portions of this preculture are transinoculated into  
10 ml of production medium (25 g/l (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 2 g/l KH<sub>2</sub>PO<sub>4</sub>,  
1 g/l MgSO<sub>4</sub>\*7H<sub>2</sub>O, 0.03 g/l FeSO<sub>4</sub>\*7H<sub>2</sub>O, 0.018 g/l MnSO<sub>4</sub>\*1H<sub>2</sub>O,  
30 g/l CaCO<sub>3</sub>, 20 g/l glucose, 50 mg/l ampicillin) and the  
batch is incubated for 48 hours at 37°C. The formation of  
30 L-threonine by the starting strain MG442 is investigated in  
the same manner, but no addition of ampicillin to the  
medium takes place. After the incubation the optical  
density (OD) of the culture suspension is determined with

an LP2W photometer from Dr. Lange (Düsseldorf, Germany) at a measurement wavelength of 660 nm.

The concentration of L-threonine formed is then determined in the sterile-filtered culture supernatant with an amino  
5 acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column reaction with ninhydrin detection.

The result of the experiment is shown in Table 1.

Table 1

Strain	OD (660 nm)	L-Threonine g/l
MG442	5.6	1.4
MG442/pTrc99A	3.8	1.3
MG442/pTrc99AsodA	3.9	2.3

10

Brief Description of the Figure:

Figure 1: Map of the plasmid pTrc99AsodA containing the sodA gene.

The length data are to be understood as approx. data. The  
15 abbreviations and designations used have the following meaning:

- Amp: Ampicillin resistance gene
- lacI: Gene for the repressor protein of the trc promoter
- 20 • Ptrc: trc promoter region, IPTG-inducible
- sodA: Coding region of the sodA gene

- 5S: 5S rRNA region
- rrnBT: rRNA terminator region

The abbreviations for the restriction enzymes have the following meaning

- 5 • BglI: Restriction endonuclease from *Bacillus globigii*  
(ATCC 49760)
- Eco91I: Restriction endonuclease from *Escherichia coli*  
RFL 91
- 10 • HindIII: Restriction endonuclease from *Haemophilus*  
influenzae
- PvuII: Restriction endonuclease from *Proteus vulgaris*  
(ATCC 13315)
- XbaI: Restriction endonuclease from *Xanthomonas*  
campestris

**What is claimed is:**

1. A process for the preparation of L-amino acids, in particular L-threonine, which comprises carrying out the following steps:
  - 5 a) fermentation of microorganisms of the Enterobacteriaceae family which produce the desired L-amino acid and in which at least the *sodA* gene, or the nucleotide sequence which codes for this, is enhanced, in particular over-expressed,
  - 10 b) concentration of the desired L-amino acid in the medium or in the cells of the microorganisms, and
  - c) isolation of the desired L-amino acid, constituents of the fermentation broth and/or the biomass in its entirety or portions (> 0 to 100%) thereof
  - 15 optionally remaining in the product.
2. A process as claimed in claim 1, wherein microorganisms in which further genes of the biosynthesis pathway of the desired L-amino acid are additionally enhanced are employed.
- 20 3. A process as claimed in claim 1, wherein microorganisms in which the metabolic pathways which reduce the formation of the desired L-amino acid are at least partly eliminated are employed.
4. A process as claimed in claim 1, wherein the expression of the polynucleotide which codes for the *sodA* gene is
- 25 increased.
5. A process as claimed in claim 1, wherein the regulatory and/or catalytic properties of the polypeptide (protein) for which the polynucleotide *sodA* codes are
- 30 improved or increased.

6. A process as claimed in claim 1, wherein, for the preparation of L-amino acids, microorganisms of the Enterobacteriaceae family in which in addition at the same time one or more of the genes chosen from the group consisting of:
- 5 6.1 the thrABC operon which codes for aspartate kinase, homoserine dehydrogenase, homoserine kinase and threonine synthase,
  - 6.2 the pyc gene which codes for pyruvate carboxylase,
  - 10 6.3 the pps gene which codes for phosphoenol pyruvate synthase,
  - 6.4 the ppc gene which codes for phosphoenol pyruvate carboxylase,
  - 15 6.5 the pntA and pntB genes which code for transhydrogenase,
  - 6.6 the rhtB gene which imparts homoserine resistance,
  - 6.7 the mgo gene which codes for malate:quinone oxidoreductase,
  - 20 6.8 the rhtC gene which imparts threonine resistance,
  - 6.9 the thrE gene which codes for the threonine export protein,
  - 25 6.10 the gdhA gene which codes for glutamate dehydrogenase,
  - 6.11 the hns gene which codes for the DNA-binding protein HLP-II,

- 6.12 the pgm gene which codes for phosphoglucomutase,
- 6.13 the fba gene which codes for fructose biphosphate aldolase,
- 5 6.14 the ptsH gene which codes for the phosphohistidine protein hexose phosphotransferase,
- 6.15 the ptsI gene which codes for enzyme I of the phosphotransferase system,
- 10 6.16 the crr gene which codes for the glucose-specific IIA component,
- 6.17 the ptsG gene which codes for the glucose-specific IIBC component,
- 6.18 the lrp gene which codes for the regulator of the leucine regulon,
- 15 6.19 the mopB gene which codes for 10 Kd chaperone,
- 6.20 the ahpC gene which codes for the small sub-unit of alkyl hydroperoxide reductase,
- 6.21 the ahpF gene which codes for the large sub-unit of alkyl hydroperoxide reductase,
- 20 6.22 the cysK gene which codes for cysteine synthase A,
- 6.23 the cysB gene which codes for the regulator of the cys regulon,
- 25 6.24 the cysJ gene which codes for the flavoprotein of NADPH sulfite reductase,
- 6.25 the cysI gene which codes for the haemoprotein of NADPH sulfite reductase,



- 6.26 the *cysH* gene which codes for adenylyl sulfate reductase,
- 6.27 the *phoB* gene which codes for the positive regulator *PhoB* of the *pho* regulon,
- 5 6.28 the *phoR* gene which codes for the sensor protein of the *pho* regulon,
- 6.29 the *phoE* gene which codes for protein E of outer cell membrane,
- 10 6.30 the *malE* gene which codes for the periplasmic binding protein of maltose transport
- 6.31 the *pykF* gene which codes for fructose-stimulated pyruvate kinase I,
- 6.32 the *talB* gene which codes for transaldolase B,
- 15 6.33 the *rseA* gene which codes for a membrane protein which acts as a negative regulator on *sigmaE* activity,
- 6.34 the *rseC* gene which codes for a global regulator of the *sigmaE* factor,
- 20 6.35 the *pfkB* gene which codes for 6-phosphofructokinase II,
- 6.36 the *sucA* gene which codes for the decarboxylase sub-unit of 2-ketoglutarate dehydrogenase,
- 25 6.37 the *sucB* gene which codes for the dihydrolipoyltranssuccinase E2 sub-unit of 2-ketoglutarate dehydrogenase,
- 6.38 the *sucC* gene which codes for the  $\beta$ -sub-unit of succinyl-CoA synthetase,

6.39 the sucD gene which codes for the  $\alpha$ -sub-unit of succinyl-CoA synthetase,


is or are enhanced, in particular over-expressed, are fermented.

5 7. A process as claimed in claim 1, wherein, for the preparation of L-amino acids, microorganisms of the Enterobacteriaceae family in which in addition at the same time one or more of the genes chosen from the group consisting of:

- 10 7.1 the tdh gene which codes for threonine dehydrogenase,
- 7.2 the mdh gene which codes for malate dehydrogenase,
- 15 7.3 the gene product of the open reading frame (orf) yjfA,
- 7.4 the gene product of the open reading frame (orf) ytfP,
- 7.5 the pckA gene which codes for phosphoenolpyruvate carboxykinase,
- 20 7.6 the poxB gene which codes for pyruvate oxidase,
- 7.7 the aceA gene which codes for isocitrate lyase,
- 7.8 the dgsA gene which codes for the DgsA regulator of the phosphotransferase system,
- 25 7.9 the fruR gene which codes for the fructose repressor,
- 7.10 the rpoS gene which codes for the sigma<sup>38</sup> factor

is or are attenuated, in particular eliminated or reduced in expression, are fermented.

Figure 1:



## SEQUENCE PROTOCOL

5     <110> Degussa AG

      <120> Process for the preparation of L-amino acids using  
          strains of the Enterobacteriaceae family which contain an  
          enhanced *sodA* gene

10    <130> 020289BT

      <160> 2

      <170> PatentIn version 3.1

15    <210> 1

      <211> 32

      <212> DNA

      <213> artificial sequence

20    <220>

      <221> Primer

      <222> (1)..(32)

      <223> *sodA1*

25    <400> 1

      cgcggcacatct agaattctgac gcccgacaat ac 32

30    <210> 2

      <211> 26

      <212> DNA

      <213> artificial sequence

35    <220>

      <221> Primer

      <222> (1)..(26)

      <223> *sodA2*

40    <400> 2

      cgcctcattg aagcttgccg caaatg 26

45

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
30 January 2003 (30.01.2003)

PCT

(10) International Publication Number  
**WO 03/008613 A3**

(51) International Patent Classification<sup>7</sup>: C12N 15/53,  
9/02, C12P 13/04, 13/08 // (C12P 13/08, C12R 1:19)

(21) International Application Number: PCT/EP02/07373

(22) International Filing Date: 3 July 2002 (03.07.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
101 35 053.8 18 July 2001 (18.07.2001) DE  
60/306,869 23 July 2001 (23.07.2001) US

(71) Applicant (for all designated States except US): DE-  
GUSSA AG [DE/DE]; Bennigsenplatz 1, 40474 Düsseldorf (DE).

(72) Inventor; and

(75) Inventor/Applicant (for US only): RIEPING, Mechthild  
[DE/DE]; Mönkebergstrasse 1, 33619 Bielefeld (DE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,

GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,  
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,  
VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),  
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),  
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,  
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,  
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,  
GW, ML, MR, NE, SN, TD, TG).

**Declaration under Rule 4.17:**

— of inventorship (Rule 4.17(iv)) for US only


**Published:**

— with international search report

(88) Date of publication of the international search report:  
18 December 2003

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROCESS FOR THE PREPARATION OF L-AMINO ACIDS USING STRAINS OF THE ENTEROBACTERIACEAE FAMILY WHICH CONTAIN AN ENHANCED SODA GENE



(57) Abstract: The invention relates to a process for the preparation of L-amino acids, in particular L-threonine, in which the following steps are carried out: a) fermentation of microorganisms of the Enterobacteriaceae family which produce the desired L-amino acid and in which at least the sodA gene, or the nucleotide sequence which codes for this, is enhanced, in particular over-expressed, b) concentration of the desired L-amino acid in the medium or in the cells of the bacteria, and c) isolation of the desired L-amino acid.



WO 03/008613 A3

# INTERNATIONAL SEARCH REPORT

International Classification No.  
PCT/EP 02/07373

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/53 C12N9/02 C12P13/04 C12P13/08 //(C12P13/08,  
C12R1:19)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE, FSTA, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 1 077 261 A (DEGUSSA ;CT NAT DE (FR)) 21 February 2001 (2001-02-21) claims 7-23; examples 3-5	1-7
Y	US 4 278 765 A (DEBABOV VLADIMIR G ET AL) 14 July 1981 (1981-07-14) cited in the application the whole document	1-7
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

11 June 2003

Date of mailing of the international search report

18/06/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Devijver, K

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 02/07373

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E, L	<p>WO 03 008605 A (RIEPING MECHTHILD ;DEGUSSA (DE)) 30 January 2003 (2003-01-30) (L: Priority) claims 1-7</p> <p>-&amp; WO 03 008615 A (RIEPING MECHTHILD ;DEGUSSA (DE)) 30 January 2003 (2003-01-30)</p> <p>-&amp; WO 03 008614 A (RIEPING MECHTHILD ;DEGUSSA (DE)) 30 January 2003 (2003-01-30)</p> <p>-&amp; WO 03 008612 A (RIEPING MECHTHILD ;DEGUSSA (DE)) 30 January 2003 (2003-01-30)</p> <p>-&amp; WO 03 008611 A (RIEPING MECHTHILD ;DEGUSSA (DE)) 30 January 2003 (2003-01-30)</p> <p>-&amp; WO 03 008610 A (RIEPING MECHTHILD ;DEGUSSA (DE)) 30 January 2003 (2003-01-30)</p> <p>-&amp; WO 03 008609 A (RIEPING MECHTHILD ;DEGUSSA (DE)) 30 January 2003 (2003-01-30)</p> <p>-&amp; WO 03 008608 A (RIEPING MECHTHILD ;DEGUSSA (DE)) 30 January 2003 (2003-01-30)</p> <p>-&amp; WO 03 008606 A (RIEPING MECHTHILD ;DEGUSSA (DE)) 30 January 2003 (2003-01-30)</p> <p>-----</p>	1-7
A	<p>TOUATI D: "CLONING AND MAPPING OF THE MANGANESE SUPEROXIDE DISMUTASE GENE SOD-A OF ESCHERICHIA-COLI K-12" JOURNAL OF BACTERIOLOGY, vol. 155, no. 3, 1983, pages 1078-1087, XP008017570 ISSN: 0021-9193 cited in the application</p> <p>-----</p>	
A	<p>WO 98 04715 A (ARCHER DANIELS MIDLAND CO) 5 February 1998 (1998-02-05)</p> <p>-----</p>	
T	<p>WO 03 008600 A (DEGUSSA ;HERMANN THOMAS (DE)) 30 January 2003 (2003-01-30) claims 1-7</p> <p>-----</p>	1-7



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/EP 02/07373

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1077261	A	21-02-2001	US 6569650 B1	27-05-2003
			AU 5188100 A	15-02-2001
			BR 0003561 A	05-06-2001
			CA 2314595 A1	13-02-2001
			CN 1284560 A	21-02-2001
			EP 1077261 A2	21-02-2001
			HU 0003279 A2	29-06-2002
			JP 2001078788 A	27-03-2001
			SK 11872000 A3	12-03-2001
			US 2002137151 A1	26-09-2002
US 4278765	A	14-07-1981	SU 875663 A1	15-09-1982
			HU 190999 B	28-12-1986
WO 03008605	A	30-01-2003	DE 10135053 A1	06-02-2003
			WO 03008605 A2	30-01-2003
			WO 03008606 A2	30-01-2003
			WO 03008607 A2	30-01-2003
			WO 03008608 A2	30-01-2003
			WO 03008609 A2	30-01-2003
			WO 03008610 A2	30-01-2003
			WO 03008611 A2	30-01-2003
			WO 03008612 A2	30-01-2003
			WO 03008613 A2	30-01-2003
			WO 03008614 A2	30-01-2003
			WO 03008615 A2	30-01-2003
WO 9804715	A	05-02-1998	AU 730102 B2	22-02-2001
			AU 3899497 A	20-02-1998
			BR 9710503 A	11-01-2000
			CN 1226931 A	25-08-1999
			EP 0917578 A1	26-05-1999
			HU 9903856 A2	28-03-2000
			JP 2000515763 T	28-11-2000
			KR 2000029691 A	25-05-2000
			NO 990362 A	26-01-1999
			NZ 334032 A	29-09-2000
			PL 331351 A1	05-07-1999
			TR 9900213 T2	21-04-1999
			WO 9804715 A1	05-02-1998
			US 5939307 A	17-08-1999
			ZA 9706803 A	19-02-1998
WO 03008600	A	30-01-2003	DE 10135051 A1	06-02-2003
			WO 03008600 A2	30-01-2003
			WO 03008602 A2	30-01-2003
			WO 03008603 A2	30-01-2003
			WO 03008604 A2	30-01-2003
			WO 03008616 A2	30-01-2003